

VERÖFFENTLICHUNGEN 2015

Abt. Pflanzenzüchtung

- Ali, M., 2015: Physiological response of German winter faba bean (*Vicia faba* L.) to drought. *Journal of Crop Improvement* 29: 319-332.
- Ali, M., 2015: Genetic Linkage Maps and Homology Study of Backcross Families of German Faba Bean (*Vicia faba* L.). *Journal of Crop Improvement*, 29: 474–490, 2015.
- Ecke, W., Kampouridis, A., Ziese-Kubon, K., and A. C. Hirsch, 2015: Identification and genetic characterization by high-throughput SNP analysis of intervarietal substitution lines of rapeseed (*Brassica napus* L.) with enhanced embryogenic potential. *Theor Appl Genet* 128: 587-603.
- Ghaouti, L., A. Schierholt, and W. Link, 2015: Effect of competition between *Vicia faba* and *Camelina sativa* as a model weed in breeding for organic conditions. *Weed Research*, accepted for publication.
- Hom, N. H., A. Schierholt, C. Möllers, and H.C. Becker, 2015: Pollen genotype effects on seed quality traits in winter oilseed rape. *Crop Science* 55: 493-500.
- Polreich, S., H.C. Becker, and B.L. Maass, 2015: Accession-specific effects of repeated harvesting edible cowpea leaves on leaf yield performance, stability and reliability – a case study from northern Tanzania. *Intern J Vegetable Science*, DOI: 10.1080/19315260.2015.1039743.
- Sallam, A., and R. Martsch, and Y. Moursi. 2015. Genetic variation in morpho-physiological traits associated with frost tolerance in faba bean (*Vicia faba* L.). *Euphytica* 205: 395-408.
- Sallam, A., and R. Martsch, 2015. Association mapping for frost tolerance using multi-parent advanced generation inter-cross (MAGIC) population in faba bean (*Vicia faba* L.). *Genetica* 143: 501-514.
- Sallam, A., Dhanapal, A. and L., Siyang, 2015: Association mapping of winter hardiness and yield traits in winter faba bean (*Vicia faba* L.). *Crop and Pasture Science*, accepted for publication.
- Teh, L.S., and C. Möllers, 2015: Genetic variation and inheritance of phytosterol and oil content in a doubled haploid population derived from the winter oilseed rape Sansibar × Oase cross. *Vorträge für Pflanzenzüchtung* 85: 37-42.
- Teh, L., and C. Möllers, 2015. Genetic variation and inheritance of phytosterol and oil content in a doubled haploid population derived from the winter oilseed rape Sansibar × Oase cross. *Theor Appl Genet*, DOI 10.1007/s00122-015-2621-y.
- Ruland, M., S. Trouchaud, M. Senbayram, and B. Horneburg, 2015. Site-specific adaptation of lentil by natural selection and its effect on drought stress. *Actas AEL* 6: 51-52.
- Webb, A., A. Cottage, T. Wood, K. Khamassi, D. Hobbs, K. Gostkiewicz, M. White, H. Khazaei, M. Ali, D. Street, G. Duc, F. Stoddard, F. Maalouf, F. Ogbonnaya, W. Link, J. Thomas, D.M. O’Sullivan. 2015. A SNP-based consensus genetic map for synteny-based trait targeting in faba bean (*Vicia faba* L.). *Plant Biotechnology Journal*, DOI 10.1111/pbi.12371.

Monographien

Moursi, Yasser, 2014: Genetic mapping of QTL controlling salt tolerance and glucosinolates in *Brassica napus* and *Brassica oleracea*. Dissertation Georg-August-Universität Göttingen. 208 S. <http://hdl.handle.net/11858/00-1735-0000-0023-9954-F>.

Teh, Li Shia, 2015: Genetic variation and inheritance of phytosterol and oil content in winter oilseed rape (*Brassica napus* L.). Dissertation Georg-August-Universität Göttingen. 154 S. <http://hdl.handle.net/11858/00-1735-0000-0022-5D9B-E>.

Miersch, Sebastian, 2015: Nitrogen efficiency in semi-dwarf and normal hybrids of oilseed rape. Dissertation Georg-August-Universität Göttingen. 112 S. <http://hdl.handle.net/11858/00-1735-0000-0022-602B-3>.